



Mouse Anti Human CD11a FITC

PRODUCT INFORMATION

CLONE: HI111
ISOTYPE: Mouse IgG1, κ
WS.No.: IV N231
CATALOG#: A6222/ A6232
CONTENTS: FITC conjugated antibody buffered in 10mM PBS (pH 7.0) with 0.05% NaN_3 and 1% BSA.

DESCRIPTION

CD11a McAb recognizes the 180KD integrin α chain associated with CD18 (95KD integrin $\beta 2$) forming heterodimer glycoprotein called the lymphocyte function antigen-1 (LFA-1, CD11a/CD18). Leucocytes express four CD11/CD18 integrins which comprise a common $\beta 2$ subunit and distinctive α chains. The other three integrins are known as CD11b/CD18, CD11c/CD18 and CD11d/CD18. LFA-1 is present on all leucocytes and is responsible chiefly for adhesion between leucocytes during immune response and also functions in transmigration of leucocytes across endothelium. The ligands for CD11a/CD18 are intercellular adhesion molecules CD54 (ICAM-1), CD102 (ICAM-2) and CD50 (ICAM-3). Patients who have mutated CD18 genes do not express cell membrane LFA-1 (or the other $\beta 2$ integrins), a condition known as leucocyte adhesion deficiency 1 (LAD-1).

PREPARATION

The monoclonal antibody is purified from ascites by protein G affinity chromatography and is conjugated with FITC under optimum conditions.

USAGE

The FITC conjugation is tested for flow cytometric analysis using $20\mu\text{l}/10^6$ cells or $100\mu\text{l}$ peripheral blood cells.

STORAGE

Storage at 4°C , should not be frozen and avoid prolonged exposure to light.

REFERENCES

Knapp, W., B. Dorken, E. P. Rieber, et al., eds. 1989. Leucocyte Typing IV: White Cell Differentiation Antigens. P: 548, 1077 Oxford University Press, New York.

Use For Research Only